

TWISTED MODULI AND SUPERSYMMETRY BREAKING

S. F. KING

*Department of Physics and Astronomy,
University of Southampton, Southampton, SO17 1BJ, U.K.
E-mail: sfk@hep.phys.soton.ac.uk*

D. A. J. RAYNER

*Dipartimento di Fisica ‘G. Galilei’, Università di Padova and INFN,
Sezione di Padova, Via Marzola 8, I-35131 Padua, Italy
E-mail: rayner@pd.infn.it*

We discuss how localized twisted moduli in type I string theory can provide a string realization of brane world supersymmetry breaking models.

String theory offers the most appealing framework for unifying all four fundamental forces together. Much recent work has been devoted to studying the problem of supersymmetry (SUSY) breaking in the context of higher-dimensional models involving parallel branes ^a. These models are inspired by the Horava-Witten setup ¹, but studied using effective field theory (EFT) techniques with an ultraviolet cutoff that is often identified with the string scale M_* . Generic models confine the MSSM fields to a 3-brane which is spatially-separated (sequestered) along a transverse extra dimension from another 3-brane where SUSY is broken by a vacuum expectation value (VEV) F . Direct couplings between sectors are assumed to arise only from the exchange of bulk fields with masses *above* the string scale M_* . After integrating out these heavy modes, we find that direct couplings (i.e. scalar masses m_ϕ) are exponentially suppressed by the separation, R

$$m_{\phi,tree}^2 \sim e^{-M_* R} \frac{F^2}{M_*^2} \quad (1)$$

Hence, SUSY breaking is (predominantly) mediated across the bulk via the superconformal anomaly ² or gaugino loops ³, so that the leading contributions to scalar masses arise at 1-loop as shown in Figure 1.

^aHere we use the term “brane” to describe any subspace of the higher-dimensional theory.




Figure 1. A generic parallel-brane setup where MSSM fields are sequestered away from SUSY-breaking along an extra dimension x_5 .

These types of models are phenomenologically appealing since they alleviate the SUSY flavour problem by suppressing flavour-changing neutral-currents (since loop corrections are flavour-blind), and also provide a simple and elegant realization of hidden-sector scenarios. In traditional 4D models, SUSY breaking originated in a hidden-sector that was only *hidden* from the visible MSSM fields by the weakness of direct couplings, e.g. Planck-scale suppressed couplings in gravity mediated models. In contrast, extra-dimensional theories provide a geometric way to separate the two sectors such that they occupy completely distinct and (apparently) unrelated physical spaces!

However, there has been recent criticism of these *sequestered* models that question whether the required form of the Kähler potential (essentially of the no-scale type) can be realized in generic string compactifications⁴. These authors dispute the original claim of Ref.² that direct couplings between sectors are suppressed, by arguing that the exchange of bulk modes with masses *below* the string scale M_* do not miraculously cancel, but instead induce inter-brane couplings that generate non-trivial scalar masses at tree-level. Nevertheless, the sequestering mechanism is still highly attractive which has motivated our attempts to embed sequestering into type I models⁵ using twisted moduli⁶.

Twisted moduli (Y_k) are closed string states that become trapped at orbifold fixed points during compactification. They play an important role in cancelling anomalies in type I theory through a generalized Green-Schwarz mechanism⁷. They appear in the gauge kinetic function at tree-level (e.g. $f_9 = S + \sum_k Y_k$ for a gauge group arising from a D9-brane) and therefore provide a new source of gaugino masses if their auxiliary F-terms acquire

non-zero VEVs ⁶. More recently, they have been shown to be crucial for dilaton stabilization in type I models using only a *single* gaugino condensate ⁸. Their localization at 4D orbifold fixed points implies that they only couple to string states that overlap with them, and so if MSSM states are localized at a distance from the twisted moduli, we anticipate a suppression of direct couplings.




Figure 2. We propose an equivalence between intersecting D-brane constructions and parallel brane EFTs, where the matter (hidden sector) brane is like open string states $C^{5_1 5_2}$ (twisted moduli Y_2) respectively.

In Figure 2, we propose that parallel brane EFTs are equivalent to string constructions involving intersecting D5-branes and twisted moduli. Matter fields are identified with open string states that stretch between the D-branes and are effectively localized at the 4D intersection point, while the hidden sector brane is like a twisted moduli localized at a fixed point away from the origin. We can also include the contributions from dilaton S and untwisted moduli T_i fields that live in the full 10D space and provide a string theoretic description of gravity mediation. In the limit that SUSY breaking originates from the twisted moduli F-term VEV, we expect that scalar masses ($m_{C^{5_1 5_2}}$) have an explicit dependence on the distance r from the SUSY breaking. However, using the standard heterotic-inspired Kähler potential for open string states ($C^{5_1 5_2}$) at the intersection between branes ⁹,

$$K_{C^{5_1 5_2}} = \frac{1}{\sqrt{S + \bar{S}} \sqrt{T_3 + \bar{T}_3}} |C^{5_1 5_2}|^2 \quad (2)$$

we discover that the masses are completely independent of this separation. This leads us to postulate a modified Kähler potential for these sequestered states

$$K'_{C^{5_1 5_2}} = \exp \left[\frac{1}{6} (1 - \eta) (Y_2 + \bar{Y}_2)^2 \right] \frac{1}{\sqrt{S + \bar{S}} \sqrt{T_3 + \bar{T}_3}} |C^{5_1 5_2}|^2 \quad (3)$$

where $\eta = e^{-M_* r}$ or $e^{-(M_* r)^2}$. Notice that we can recover the standard result in Eq. 2 by taking the limit that $r \rightarrow 0$. We attribute the suppression to either (i) the propagation of heavy modes with masses above the string scale ^{2,3} ($\eta = e^{-M_* r}$); or (ii) arising from non-perturbative worldsheet instanton corrections ¹⁰ that are interpreted as (massive) strings stretched between the different fixed points ($\eta = e^{-(M_* r)^2}$).

Using our modified Kähler potential, we use a model-independent parametrization of F-terms involving Goldstino angles ^{9,11} to derive general expressions for trilinears and scalar masses, where different limits of these Goldstino angles correspond to dilaton (S), moduli (T_i) or twisted moduli (Y_k) dominated SUSY breaking. We anticipate that our general expressions can be used to extend previous type I theory parameter scans ¹², but incorporating both gravity and gaugino mediated contributions.

References

1. P. Horava and E. Witten, Nucl. Phys. B **460** (1996) 506 [arXiv:hep-th/9510209]; E. Witten, Nucl. Phys. B **471** (1996) 135 [arXiv:hep-th/9602070]; P. Horava and E. Witten, Nucl. Phys. B **475** (1996) 94 [arXiv:hep-th/9603142].
2. L. Randall and R. Sundrum, Nucl. Phys. B **557** (1999) 79 [arXiv:hep-th/9810155].
3. D. E. Kaplan, G. D. Kribs and M. Schmaltz, Phys. Rev. D **62** (2000) 035010 [arXiv:hep-ph/9911293]; Z. Chacko, M. A. Luty, A. E. Nelson and E. Ponton, JHEP **0001** (2000) 003 [arXiv:hep-ph/9911323].
4. A. Anisimov, M. Dine, M. Graesser and S. Thomas, Phys. Rev. D **65** (2002) 105011 [arXiv:hep-th/0111235]; arXiv:hep-th/0201256; C. P. Burgess, E. Filotas, M. Klein and F. Quevedo, arXiv:hep-th/0209190; Also see M. Dine's talk in these proceedings, arXiv:hep-ph/0210255.
5. S. F. King and D. A. J. Rayner, JHEP **0207** (2002) 047 [arXiv:hep-ph/0111333]; Nucl. Phys. B **607** (2001) 77 [arXiv:hep-ph/0012076].
6. K. Benakli, Phys. Lett. B **475** (2000) 77 [arXiv:hep-ph/9911517].
7. L. E. Ibanez, R. Rabadan and A. M. Uranga, Nucl. Phys. B **542** (1999) 112 [arXiv:hep-th/9808139].
8. S. A. Abel and G. Servant, Nucl. Phys. B **597** (2001) 3 [arXiv:hep-th/0009089]; Nucl. Phys. B **611** (2001) 43 [arXiv:hep-ph/0105262].
9. L. E. Ibanez, C. Munoz and S. Rigolin, Nucl. Phys. B **553** (1999) 43 [arXiv:hep-ph/9812397].
10. S. Hamidi and C. Vafa, Nucl. Phys. B **279** (1987) 465; L. J. Dixon, D. Friedan, E. J. Martinec and S. H. Shenker, Nucl. Phys. B **282** (1987) 13; L. E. Ibanez, Phys. Lett. B **181** (1986) 269.
11. A. Brignole, L. E. Ibanez and C. Munoz, Nucl. Phys. B **422** (1994) 125 [Erratum-ibid. B **436** (1995) 747] [arXiv:hep-ph/9308271]; A. Brignole, L. E. Ibanez, C. Munoz and C. Scheich, Z. Phys. C **74** (1997) 157 [arXiv:hep-ph/9508258].
12. S. A. Abel, B. C. Allanach, L. E. Ibanez, M. Klein and F. Quevedo, JHEP

0012 (2000) 026 [arXiv:hep-ph/0005260]; B. C. Allanach, D. Grellscheid and F. Quevedo, JHEP **0205** (2002) 048 [arXiv:hep-ph/0111057];